

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/141314>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

The catalytic mechanism of aromatic nitration by cytochrome P450 TxtE: Involvement of a ferric-peroxynitrite intermediate

Savvas Louka,^{§,⊥,‡} Sarah M. Barry,^{#,‡,†} Derren J. Heyes,^{§,&} M. Qadri E. Mubarak,^{§,⊥} Hafiz Saqib Ali,^{§,&} Lona Alkhalaf,[#] Andrew W. Munro,^{§,&} Nigel S. Scrutton,^{*,§,&} Gregory L. Challis^{*,§,#,‡} and Sam P. de Visser^{*,§,⊥}

[§] The Manchester Institute of Biotechnology, The University of Manchester, 131 Princess Street, Manchester M1 7DN, United Kingdom

[⊥] Department of Chemical Engineering and Analytical Science, The University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom

[&] Department of Chemistry, The University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom

[#] Department of Chemistry, University of Warwick, Coventry CV4 7AL, United Kingdom

[§] Department of Biochemistry and Molecular Biology, Monash University, Clayton VIC 3800, Australia

^{*} ARC Centre for Excellence in Innovations in Peptide and Protein Science, Monash University, Clayton, VIC 3800, Australia

ABSTRACT: The cytochromes P450 are heme-dependent enzymes that catalyze many vital reaction processes in the human body related to biodegradation and biosynthesis. They typically act as mono-oxygenases; however, the recently discovered P450 subfamily TxtE utilizes O₂ and NO to nitrate aromatic substrates such as L-tryptophan. A direct and selective aromatic nitration reaction may be useful in biotechnology for the synthesis of drugs or small molecules. Details of the catalytic mechanism are unknown and it has been suggested that the reaction should proceed through either an iron(III)-superoxo or an iron(II)-nitrosyl intermediate. To resolve this controversy, we used stopped-flow kinetics to provide evidence for a catalytic cycle where dioxygen binds prior to NO to generate an active iron(III)-peroxynitrite species that is able to nitrate L-Trp efficiently. We show that the rate of binding of O₂ is faster than that of NO and also leads to L-Trp nitration, while little evidence of product formation is observed from the iron(II)-nitrosyl complex. To support the experimental studies we performed density functional theory studies on large active site cluster models. The studies suggest a mechanism involving an iron(III)-peroxynitrite that splits homolytically to form an iron(IV)-oxo heme (Compound II) and a free NO₂ radical via a small free energy of activation. The latter activates the substrate on the aromatic ring, while Compound II picks up the *ipso*-hydrogen to form the product. The calculations give small reaction barriers for most steps in the catalytic cycle and, therefore, predict fast product formation after the formation of an iron(III)-peroxynitrite complex. These findings provide the first detailed insight into the mechanism of nitration by a member of the TxtE subfamily and highlight how the enzyme facilitates this novel reaction chemistry.

INTRODUCTION.

The cytochromes P450 (P450s or CYPs) are vital enzymes for human health with, e.g., key functions in the liver related to the metabolism of drugs and xenobiotics; and as such their activity and reactivity is of interest to drug delivery and development.^{1–5} In addition, they also catalyze key steps in the biosynthesis of natural products and hormones, including estrogen.^{6–10} Their diverse reactivities and ability to activate inert C–H bonds have resulted in the P450s gaining significant interest due to their potential as

biocatalysts in the synthesis of high value chemicals.^{11–13} However, factors including requirement for expensive redox partners, as well as a lack of knowledge of many facets of their structure and activity, has limited their applications. The P450s contain a heme prosthetic group that is typically linked to the protein via a proximal coordination to the heme iron through a cysteinate residue.^{14–16} The P450s typically react as mono-oxygenases, where they transfer one atom of molecular oxygen to a substrate, while the second oxygen atom of O₂ is reduced to water.^{1–4,17–19}




Figure 1. (a) Active site structure of P450 TxtE as taken from the 4TPO pdb file. (b) Proposed reaction mechanism of dioxygen and NO activation of the heme in TxtE and possible reactive products considered here. (c) Overall nitration reaction catalyzed by P450 TxtE.

In general, the P450s, can catalyze diverse reactions, including aliphatic or aromatic hydroxylation, epoxidation and sulfoxidation, in addition to pathways leading to desaturation, C–C bond cleavage and O-dealkylation.^{20,21} P450 isozymes show substantial differences in catalytic properties and the shape and size of the substrate binding pocket. Consequently, many questions remain related to their catalytic cycle and substrate activation processes.

Recently, two novel P450 subfamilies were discovered that are able to regiospecifically nitrate aromatic amino acids including L-tryptophan and L-tyrosine.^{22–31} The P450 TxtE isozyme activates L-tryptophan at the 4-position, while the P450 RuFO isozyme binds L-tyrosine and adds an NO₂ group to the 3-position. In actinobacteria, TxtE catalyzed nitration is part of the biosynthetic pathway of the phytotoxin thaxtomin in *Streptomyces scabies*, while tyrosine nitration catalyzed by RuFO is a key step in the biosynthesis of rufomycin, a natural product peptide antibiotic with activity against *Mycobacterium tuberculosis*. Furthermore, the L-Phe residue in the resulting N,N-dimethyldiketopiperazine is hydroxylated at the C^α and aromatic ring positions by the bifunctional P450 TxtC.^{22–24,32–34} The substrate nitration reactions are not restricted to one class of P450 isozymes, as addition of peroxynitrite to the monooxygenase P450 3A4 was found to lead to tyrosine nitration efficiently.³⁵ Direct and selective aromatic nitration is difficult to achieve chemically and thus enzyme catalyzed processes have significant potential in biotechnology and therefore we decided to investigate the catalytic mechanism of the TxtE P450 enzyme using a combined experimental and computational approach.

A TxtE crystal structure was reported with the substrate L-Trp bound,²⁴ and an extract of the active site is shown in

Figure 1a. The heme iron is bound to the thiolate group of Cys₃₅₇, the proximal axial ligand, and the L-Trp substrate is positioned in the active site in a tight orientation with hydrogen bonding interactions to several amino acids, including Arg₅₉ (Figure 1). Unfortunately, due to their transient nature, few catalytic cycle intermediates have been identified and characterized for P450 TxtE. However, it has been established that the enzyme binds L-tryptophan in the active site and utilizes O₂ and NO to convert the substrate to L-4-nitrotryptophan selectively, using external redox partners for provision of electrons.^{22–24} Free superoxide (O₂^{-•}) and NO in a water environment react within the diffusion limit to form peroxynitrite efficiently.³⁶ However, on a metal center the order of binding of NO versus O₂ determines the reaction products and the formation of nitrate, nitrite or peroxynitrite, which all react differently with substrates.³⁶ A catalytic cycle of TxtE was hypothesized that proceeds via an iron(III)-peroxynitrite intermediate.²² Previous studies on myoglobin-bound peroxynitrite showed that with open substrate-binding pockets and access of protons, the peroxynitrite was quickly converted to nitrate and no aromatic nitration process was obtained.³⁷

Little is known about the catalytic cycle of P450 TxtE and only the ferric-H₂O resting state (Figure 1a) and the water-displaced ferric states have been characterized.^{22–24} A P450 TxtE catalytic cycle has been proposed where substrate binding displaces the water molecule from the resting state and, after heme iron reduction, is followed by dioxygen binding to form an iron(II)-dioxygen or an iron(III)-superoxo intermediate, which subsequently reacts with NO to form a ferric-peroxynitrite intermediate.²² The ferric-peroxynitrite complex could react directly with substrate or either homolytically cleave to form an iron(IV)-

oxo species called Compound II (CpdII) and NO_2^\bullet , or heterolytically cleave to give an iron(IV)-oxo heme cation radical called Compound I and NO_2^- . This would give the P450 nitration isozymes a distinctive catalytic cycle that deviates from the P450 mono-oxygenases, wherein the ferric-superoxo form is reduced and doubly protonated to form CpdI and a water molecule.^{16–18,38–42} Spectroscopic and kinetic studies characterized CpdI as the active species in the *Pseudomonas putida* P450 CAM enzyme.⁴³ Although the one-electron reduced form of CpdI, i.e. CpdII, is not part of the mono-oxygenase catalytic cycle, it can be formed upon reduction of CpdI.

Another heme-dependent enzyme, namely nitric oxide synthase (NOS) has also been shown to activate peroxynitrite.^{44,45} However, in the NOS enzymes, addition of excess peroxynitrite led to the production of the ferrous-nitrosyl species instead. A similar observation was made for P450 BM₃ where excess peroxynitrite (ONOO^-) yielded iron(II)-NO products rather than the anticipated CpdI or CpdII products.⁴⁶ This is in contrast to work of Groves et al⁴⁷, who added NO to oxygenated myoglobin to detect a ferryl-oxo species, i.e. CpdII, and free NO_2^\bullet .⁴⁷ In particular, the reaction of the iron(III)-superoxo species with NO was found to yield peroxynitrite rapidly. The order by which dioxygen and NO are added to the protein may have an effect on whether ferrous-nitrosyl, CpdI or CpdII is formed, and therefore we decided to study the early stages of the catalytic cycle of TxtE, and particularly focused on the formation of the iron(II)-NO and ferric-peroxynitrite species.

In this study we investigated the order of O_2 and NO binding in TxtE, and whether free NO_2^\bullet or NO_2^- is generated and is involved in the catalytic mechanism of L-Trp nitration by P450 TxtE. In addition, our work focused on understanding the selectivity of substrate nitration and shows that an initial precursor iron(III)-peroxynitrite complex homolytically cleaves into an iron(IV)-oxo heme and NO_2^\bullet . As substrate, L-tryptophan is located close to the formed NO_2^\bullet and it is activated rapidly to form nitrated products. Clearly, substrate binding must be a precursor to NO addition to an iron(III)-superoxo intermediate in the catalytic cycle. Our work rationalizes substrate nitration by P450 isozymes and gives insights into how to further engineer P450 enzymes for more efficient biotechnological applications.

METHODS.

Experiment.

Sample preparation. N-terminal His-tagged TxtE protein was expressed and purified based on a previously described protocol.²² For large-scale protein production 5×1 L cultures were grown at 37 °C to an OD_{600} of ~0.6 and induced overnight at 15 °C by the addition of 1 mM isopropyl β -D-thiogalactopyranoside. Cells were harvested by centrifugation and the pellet was resuspended in 100 mL buffer B₁ (20 mM Tris-HCl buffer, pH 8.0, 100 mM NaCl, 20 mM imidazole, 1 mM DTT and 10% glycerol). The cells were lysed in

the presence of 1 mM phenylmethylsulfonyl fluoride using a French Press (17,000 psi internal cell pressure) followed by sonication for 2 min (Ultrasonic processor, Jencons). After removal of the cellular debris by centrifugation ($18,000 \times g$ for 20 min at 4 °C), the supernatant was applied to a 5 mL HiTrapTM HP affinity column (Nickel Sepharose High Performance, GE Healthcare) equilibrated with buffer B₁ (flow rate 2 mL min⁻¹). Unbound proteins were removed by washing with 30 mL buffer B₁ and the His₆-TxtE fusion protein was eluted with ~10 mL of buffer B₂ (25 mM Tris-HCl, pH 8.0, 100 mM NaCl, 200 mM imidazole, 1 mM DTT, 10% glycerol). The protein was further purified by gel filtration using 110 mL Superdex 12 prep grade resin (GE Healthcare) equilibrated with buffer B₃ (25 mM Tris-HCl, pH 8.0, 100 mM NaCl, 1 mM DTT and 10% glycerol). The sample was eluted with ~64 mL of buffer B₃ (flow rate 0.75 mL min⁻¹, column dead volume 40 mL) and fractions containing His₆-TxtE were analyzed by electrophoresis (10% SDS-PAGE, Figure S1, Supporting Information), pooled and concentrated using an Amicon[®] Ultracentrifugal filtration device with a 30,000 molecular weight cutoff membrane (Millipore). Pure His₆-TxtE was buffer exchanged into buffer B₄ (25 mM Tris-HCl, pH 8.0, 100 mM NaCl, and 30% glycerol), aliquoted and frozen at -80 °C. The purified protein was confirmed to be active using the previously described assay.²² UV-Vis absorbance spectra were measured using a Cary 50 UV-Vis spectrophotometer (Agilent Technologies).

Stopped-flow spectroscopy. Rapid kinetic measurements were carried out using an SX20 rapid mixing stopped-flow spectrophotometer (Applied Photophysics Ltd, Leatherhead) placed inside a Belle Technology anaerobic chamber (oxygen levels < 2 ppm) as previously described.⁴⁸ Proteins and substrates were dissolved in buffer (25 mM Tris, pH 8.0) and degassed, unless otherwise stated. L-tryptophan (0.5 mM) and sodium dithionite (15 mM) were added to TxtE protein (150 μM) to reduce it to the high-spin iron(II) form. To remove excess dithionite, the protein was passed through a PD10 desalting column equilibrated with 20 mM Tris pH 8 containing 0.15 mM L-Trp (which was kept in the glovebox overnight prior to all measurements and to ensure removal of all oxygen traces) and eluted in a volume of ~5 mL. To confirm that the protein heme iron was reduced and high-spin (L-Trp was bound), a sample was analyzed by UV-vis spectroscopy in a sealed cuvette. The protein was maintained in a reduced, high-spin state for several hours. All stopped-flow experiments were carried out at 4 °C using a final concentration of 2 μM TxtE and 75 μM L-Trp. For NO experiments with degassed buffer, the sample was bubbled with NO for 10 minutes to saturate the buffer. To ascertain that the NO did not contain impurities such as small traces of NO_2 that could affect the results,^{36,49–52} some experiments were repeated with purified NO that had first been passed through a 10% (w/v) solution of Ascarite (Sigma Aldrich) to remove any potential impurities caused by disproportionation of the NO gas from the cyl-

inder. None of these studies gave significantly different results from the experiments with non-purified NO, see Supporting Information Figure S5.

For O₂ experiments a sealed tube containing air-saturated buffer (~200 μ M O₂) was transferred to the anaerobic chamber. Serial dilutions of each solution were made just prior to use by using degassed buffer as required. Time-resolved spectral changes were recorded using a photodiode array detector and single wavelength kinetic transients were measured at 435 nm (for O₂) or 445 nm (for NO) to determine the rates of gas binding. The rates of NO and O₂ binding were measured in single mixing stopped-flow experiments (experimental set up shown schematically in Figure S2, Supporting Information) over a range of concentrations. Double mixing stopped-flow experiments were carried out to determine the order of gas binding to the protein (experimental set up shown schematically in Figure S3, Supporting Information).

The samples from the double mixing stopped-flow experiments were collected after each mix and frozen for further analysis by LC-HRMS on a Bruker MicroTOF mass spectrometer. LC-HRMS conditions for analysis of 4-L-nitrotryptophan production were as follows: LC-HRMS was performed on a Sigma Ascentis Express column (C18, 150 \times 2.1 mm, 2.7 μ m) attached to a Dionex 3000RS UHPLC coupled with a Bruker MaXis Q-TOF mass spectrometer. The mobile phase A was water containing 0.1% formic acid and mobile phase B was methanol containing 0.1% formic acid. The flow rate was set at 0.2 mL min⁻¹ and the absorbance was measured at 220 nm. The following elution profile was used: 0–5 min – 100% A; 5–17.4 min – 100% A to 100% B; 17.4–22.4 min – 100% B; then equilibrate back to 100% A.


The mass spectrometer was operated in electrospray positive mode with a scan range of 50–2,000 *m/z*. The source conditions were: end plate offset at –500 V; capillary at –4500 V; nebulizer gas (N₂) at 1.6 bar; dry gas (N₂) at 8 L min⁻¹; dry temperature at 180°C. Ion transfer conditions were: ion funnel RF at 200 V_{pp}; multiple RF at 200 V_{pp}; quadrupole low mass at 55 *m/z*; collision energy at 5.0 eV; collision RF at 600 V_{pp}; ion cooler RF at 50–350 V_{pp}; transfer time was set to 121 μ s; pre-Pulse storage time was set to 1 μ s. Calibration was carried out with sodium formate (10 mM) through a loop injection of 20 μ L of standard solution at the beginning of each run.

Theory.

Density functional theory calculations were performed on active site model complexes of TxtE based on the crystal structure coordinates of the 4TPO protein data-bank (pdb) file.²⁴ We followed set-up procedures as discussed and described previously, but we will summarize the key facts here.^{53,54} Our initial calculations used a minimal active site model (model A) that included an iron(III)-peroxynitrite-heme (with side chains replaced by hydrogen atoms) and thiolate for the axial cysteinate ligand. A complete trypto-

phan amino acid was included as substrate, where the carboxylic acid group was deprotonated and the amino group protonated. Hence, the overall charge of model A was –1 and the system was calculated in the individual doublet and quartet spin states. No constraints were put on this model.

A second and more elaborate model (model B) was subsequently studied that included the substrate binding pocket as obtained from the 4TPO pdb file with key hydrogen bonding interactions and salt bridges between substrate and protein (Scheme 1).²⁴ Thus, model B contains model A with one propionate side chain of the heme included and was expanded with part of the substrate binding pocket, namely the Arg₅₉ and Tyr₈₉ side chains, the small peptide chains Asn₂₉₃–Gln₂₉₄–Phe₂₉₅–Thr₂₉₆–Trp₂₉₇–Arg₂₉₈, and Glu₃₉₄–Phe₃₉₅, whereby the Gln₂₉₄ residue was abbreviated to Gly. In the starting structure, the Trp substrate in Model B had a zwitterionic state with the amino group protonated and the carboxylate group deprotonated and forming a salt bridge with Arg₅₉. This model has 250 atoms and an overall charge of –1, and is shown in Scheme 1. To make sure that the structure did not deviate too much from the crystal structure coordinates, some geometric constraints were placed on several α -carbon atoms of the protein backbone in the model as identified with a star in Scheme 1. A comparison of starting and final structures of the geometry optimizations showed little changes of the protein part of the model and hence it kept the features of the protein structure intact. In the following section we will identify the model (A or B) after the label of the structure.



Scheme 1. Computational models investigated in this work. Wavy lines show where a covalent bond was broken.




Figure 2. Stopped-flow analysis of the mechanism of gas binding in TxtE. (a) Absorbance spectra measured after 50 ms using stopped-flow photodiode array measurements upon mixing of TxtE iron(II) with either anaerobic buffer, 20 μM O₂ or 20 μM NO. (b) Dependence of the rate constant for gas binding to iron(II) heme TxtE on the concentration of either O₂ or NO. The data were fitted to a straight line to determine the 2nd order rate constant for O₂ or NO binding. (c) Time-resolved spectra at selected time points from double mixing stopped-flow experiments, where iron(II) heme TxtE was first mixed with 20 μM O₂, aged for 50 ms and then mixed with 20 μM NO prior to data collection. The iron(III)-superoxo species at 425 nm is rapidly converted to the water-ligated iron(III) form at 417 nm over 100 ms. (d) Measurable amounts of the L-4-nitrotryptophan product formed in samples collected from double mixing stopped-flow experiments, where iron(II) heme TxtE was first mixed with either 20 μM O₂ or 20 μM NO, aged for 50 ms and then mixed with 100 μM NO or O₂, respectively. All raw data can be found in the Supporting Information.

For Model A, a range of possible reaction pathways for substrate nitration were explored using Gaussian-09,⁵⁵ see Supporting Information. Using the obtained insight, we then investigated the low energy pathways with the expanded cluster model B. Based on our previous experience with heme systems, we used well-tested and validated approaches that use density functional theory (DFT) meth-

ods.^{56–58} All geometries were fully optimized without constraints and transition states were characterized with a single imaginary frequency for the correct mode. For a number of transition state geometries, intrinsic reaction coordinate scans were also performed that connected them to the two local minima adjacent to the transition state (see Supporting Information).

The hybrid density functional method UB3LYP^{59,60} was used for all calculations in combination with an LACVP basis set on iron with core potential and 6-31G* on the rest of the atoms (C, N, O, H, S), basis set BS1.^{61,62} All optimizations, frequencies and geometry scans included a continuum polarized conductor model (CPCM) with a dielectric constant mimicking chlorobenzene.⁶³ To obtain more accurate energies, we did single point calculations with an LACV3P+ basis set on iron with electron core potential and 6-311+G* on the rest of the atoms: basis set BS2. These methods and approaches were used previously and shown to predict the correct regio- and chemoselectivities of reaction mechanisms, and to give free energies of activation to within 3 kcal mol⁻¹ of the experiment.⁶⁴⁻⁶⁶

For a selection of small model complexes we tested the effect of an external electric field on the binding energy of NO₂ to an indole group using the keyword "Field" as implemented in Gaussian-09.⁵⁵

RESULTS.

Stopped-flow analyses. Stopped-flow spectroscopy experiments were performed to determine the order of gas binding. The P450 catalytic cycle for a mono-oxygenation reaction starts with a substrate binding in the active site close to the iron(III)-heme in its resting state, followed by release of water from the sixth ligand position of the heme and resulting in conversion of iron from low spin (LS) to high spin (HS). This is followed by a single electron reduction of the heme from HS iron(III) to iron(II) using a redox partner, enabling molecular oxygen binding to form the iron(III)-superoxo.^{1-3,17-19,67,68} A further reduction and two protonation steps result in the formation of Compound I (CpdI), an iron(IV)-oxo heme cation radical species that is the active oxidant in oxygen atom transfer reactions. Trp nitration by P450 TxtE is believed to deviate from this mechanism and upon formation of the iron(III)-superoxo species the enzyme reacts with NO to form an iron(III)-peroxynitrite intermediate, or alternatively gives a ferric-nitrosyl complex. Indeed, in all systems studied to date the P450 TxtE nitration enzyme is found to be linked to a nitric oxide synthase enzyme that provides NO for the process by reacting L-Arg with O₂ on the heme iron.^{22,24}


In order to confirm the mechanism for TxtE and the order of O₂ and NO binding, we have used stopped-flow rapid mixing measurements to determine the order of gas binding to the protein. Initial stopped-flow measurements involved mixing the TxtE enzyme against either NO (purified or unpurified) or O₂ and showed that characteristic transient iron(II)-NO and iron(III)-superoxo species were formed in each case. Time-resolved spectra, recorded using a photodiode array detector, showed that the absorbance maximum of the heme-iron state transiently shifts from approximately 410 nm to 425 nm upon binding of O₂, and to 435 nm upon binding of NO (Figure 2a) and similar to reported literature.^{69,70} Identical spectral changes were observed when purified NO was used in the measurements,

suggesting that any effects from possible impurities caused by disproportionation of the NO source were minimal (see Figure S5, Supporting Information).

Previous studies on dioxygen binding to the P450 enzymes also showed a Soret band at 425 nm for the substrate-bound ferric-superoxo species, in agreement with the TxtE data seen here.⁷¹ Consequently, the rate of gas binding was measured at the wavelength where the maximum increase in absorbance occurred: in our case we measured the absorbance changes at 435 nm for O₂ and at 445 nm for NO over a range of O₂ and NO concentrations (Figure S4 and S5 for purified NO, Supporting Information) rather than at the Soret maximum absorbances. In each case, the rate of binding was linearly dependent on the gas concentration, yielding 2nd order rate constants of $4.43 \pm 0.03 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ for O₂ binding and $1.82 \pm 0.03 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ for NO binding (Figure 2b). Hence, the rate of binding of O₂ is approximately 2.5 times faster than that for NO, implying that O₂ is most likely to bind to the enzyme prior to NO. Moreover, this is likely to be particularly evident inside the cell where the concentration of O₂ would be expected to be much higher than that of NO.

Next, we attempted to isolate the putative iron(III)-peroxynitrite intermediate and to confirm the order of gas binding in TxtE through a double mixing stopped-flow experiment. To this end, TxtE was first mixed with O₂ or NO, allowed to age for 50 ms to form the initial iron(III)-superoxo or iron(II)-NO species, and then mixed with the second gas prior to collection of the time-resolved spectral changes. Measurements where TxtE was first mixed with NO, followed by a second mix with O₂, only appeared to show the formation of the iron(II)-NO species at 435 nm on timescales faster than 1 second (Figure S8, Supporting Information). On longer timescales (>100 s) the 435 nm peak slowly disappeared and a new peak at 425 nm was formed, which is likely to represent the iron(III)-superoxo species as the bound NO is gradually displaced by O₂ (Figure S8, Supporting Information). Hence, it appears that O₂ is unable to react directly with the iron(II)-NO intermediate to form any new species, suggesting that dioxygen must bind prior to NO in order to produce a catalytically active species.

The reverse experiments, involving an initial mix with O₂ followed by a second mix with NO, were slightly more complicated as the iron(III)-superoxo species decays over approximately 1 second, even upon mixing with anaerobic buffer. It is converted to a new species with an absorbance maximum at 417 nm, which represents the water-ligated iron(III) form of the enzyme (Figure S9, Supporting Information). Nevertheless, clear differences are observed upon binding of NO to the iron(III)-superoxo, implying that binding of dioxygen prior to NO appears to give a more likely catalytic intermediate than the reverse ordering.




Scheme 2. Reaction mechanism of TxtE as derived from stopped-flow measurements.

The iron(III)-superoxo species is converted to the same water-ligated iron(III) species much more rapidly (<100 ms) upon mixing with NO (Figure 2c and Figure S10, Supporting Information), presumably as a result of catalysis to form the L-4-nitrotryptophan product. On longer time-scales (>4s) the L-Trp substrate rebinds to this iron (III) to give rise to starting substrate-bound heme spectrum at 390 nm (Figure S8).


Further confirmation of the binding order was provided by additional double mixing experiments, where equal concentrations of O₂ and NO were initially mixed and allowed to age for 50 ms, prior to mixing with the enzyme (Figure S9). In this case, there was no observable formation of the iron(II)-NO species at 435 nm. Instead, rapid formation of the iron(III)-superoxo species was observed at 425 nm, followed by the rapid decay to the water-ligated iron(III) species at 417 nm, in a similar manner to the previous binding of NO to the iron(III)-superoxo complex. It was not possible to identify any additional catalytic intermediates in the double mixing photodiode array experiments, which implies that the rate of formation of the proposed peroxynitrite species is likely to be much slower than the subsequent reaction chemistry.

Consequently, in order to provide further evidence that the catalytic iron(III)-peroxynitrite intermediate is formed directly upon NO binding to the iron(III)-superoxo species, we collected samples from the double mixing stopped-flow experiments and measured the amount of L-4-nitrotryptophan product that had been formed (Figure 2d and Figure S10, Supporting Information). Upon mixing of the iron(II)-NO complex with different concentrations of O₂, only a small amount of L-4-nitrotryptophan was formed, presumably because the NO is gradually displaced by O₂ prior to collecting and freezing the samples from the stopped-flow instrument. However, the amount of the L-4-nitrotryptophan product is significantly higher when the iron(III)-superoxo species is mixed with NO (Figure 2d). Moreover, the amount of product increases with the con-

centration of NO used in the second mix (Figure S10, Supporting Information). This indicates that efficient reaction chemistry only occurs when O₂ binds first to form the iron(III)-superoxo species, followed by NO binding to generate the catalytically relevant iron(III)-peroxynitrite intermediate. The experimentally derived catalytic cycle of P450 TxtE, based on the stopped-flow experiments presented here, is shown in Scheme 2. The work shows that both dioxygen and NO can bind to an iron(II) pentacoordinated heme structure, although the dioxygen binds faster. Subsequently, the iron(III)-superoxo species rapidly reacts with NO and one molecule of substrate to form 4-nitro-L-tryptophan, and returns the enzyme to the resting state. Unfortunately, the peroxynitrite complex was too short-lived to characterize and study in detail, and hence a computational study was performed.





Scheme 3. DFT calculated mechanisms for iron(III)-peroxynitrite formation.



Scheme 4. Mechanistic pathways explored for the reaction of iron(III)-peroxynitrite with L-Trp on the C⁴-position using DFT methods.

Computation. To gain further insight into the catalytic cycle of TxtE and related nitrating enzymes, we pursued a computational study using model complexes based on the active site and substrate binding environment of the enzyme. Our cluster model had 250 atoms and is shown above in Scheme 1. Our initial focus was on the potential role of peroxynitrite in the reaction mechanism, but the substrate activation reactions were also investigated. As we propose free NO₂ to be formed during the catalytic cycle of TxtE, we decided to explore substrate activation on the C⁴, C⁵, C⁶ and C⁷-positions of L-Trp. In particular, the reaction energy with zero-point energy (ZPE) correction ($\Delta E + ZPE_R$) for the formation of iron(III)-peroxynitrite from either iron(III)-superoxo + NO (Eq 1) or iron(II)-nitrosyl + O₂ (Eq 2) were calculated. To this end, the individual molecules in reactions 1 and 2 were calculated with the small model complex **A** in the absence of substrate and the energy differences for these reactions are shown in Scheme 3.



Both reactions are exothermic, with values of $-11.4 \text{ kcal mol}^{-1}$ for reaction 1 and $-36.0 \text{ kcal mol}^{-1}$ for reaction 2. Note

that with the large model complex, the energy for reaction 1 was calculated to be $\Delta E + ZPE_R = -12.3 \text{ kcal mol}^{-1}$; hence the protein environment has little effect on the iron(III)-peroxynitrite bond formation.

However, the reaction rates for the two processes give the opposite ordering and a barrier height with magnitude $\Delta E^\ddagger + ZPE = 14.0 \text{ kcal mol}^{-1}$ was calculated for reaction 1, while reaction 2 is estimated to be well over 27 kcal mol^{-1} using gas-phase model complexes. This reaction barrier is in good quantitative agreement with calculations on a biomimetic iron-porphyrin model complex with axial imidazole group for an O-bound iron(III)-peroxynitrite dissociation into an iron(IV)-oxo and NO₂.⁷² Furthermore, DFT calculations of large cluster models of heme-NO and heme-O₂ species showed that O₂ binds more strongly than NO.⁷³ It should be mentioned here that the reaction of iron(III)-superoxo with NO gives an O-bound iron(III)-peroxynitrite structure, whereas the reaction of iron(II)-NO with O₂ gives the less stable N-bound iron(III)-peroxynitrite that is rapidly released from the iron center and forms a pentacoordinated iron(II)-heme and a free peroxynitrite ion. As such, the DFT calculations on small cluster models support a faster reaction of iron(III)-superoxo with NO than iron(II)-NO with O₂.




Figure 3. Potential energy landscape (with values in kcal mol⁻¹) of L-Trp nitration at the C⁴-position via exo-attack in an active site model complex B. All geometries optimized in Gaussian at UB₃LYP/BS1. The two energetic values represent $\Delta E^{\ddagger} + ZPE$ (ΔG) data with energies obtained using basis set BS2 and with ZPE and solvent corrections; free energies are given in parenthesis at 298 K and contain ZPE, thermal, solvent and entropic corrections. Optimized geometries of the transition states give bond lengths in angstroms and the imaginary frequency in cm⁻¹.

Moreover, they show that iron(II)-NO with dioxygen does not give an iron(III)-peroxynitrite directly, but instead a relatively unstable N-bound iron(III)-peroxynitrite that either reacts to form iron(III)-superoxo by NO release, or reorients to form the O-bound iron(III)-peroxynitrite complex. The driving force for converting the iron(II)-NO + O₂ into an iron(III)-superoxo and a NO molecule was calculated to be exothermic by 24.6 kcal mol⁻¹.

Next, we explored the reaction mechanism of substrate activation in TxtE, leading to substrate nitration and hydroxylation as displayed in Scheme 4. Initially, the work focused on the steps from the iron(III)-peroxynitrite complexes in the doublet and quartet spin states, and we investigated three major reaction pathways. These iron(III)-peroxynitrite complexes (^{4,2}Re) were geometry optimized with a large active site model that includes the substrate and the binding pocket. Pathways 1 and 2 start with a homolytic O–O bond cleavage in the peroxynitrite group to form Compound II, i.e. an iron(IV)-oxo heme and NO₂

(complex **Int1**) via a transition state **TS1**. Thereafter, pathways 1 and 2 diverge and substrate attack of either NO₂ or iron(IV)-oxo were investigated. In pathway 1, the free NO₂ radical attacks the substrate directly to form intermediate **Int4** via transition state **TS4**, which after hydrogen atom transfer to the iron(IV)-oxo species via transition state **TS5** leads to nitrated Trp and the ferric-hydroxide complex, i.e. protonated Compound II (**Int5**).

In pathway 2, by contrast, an electrophilic addition of the oxo to a carbon atom of the Trp substrate takes place to form a radical intermediate **Int2** via a transition state **TS2**. This is then followed by abstraction of the *ipso*-proton from the Trp carbon atom by the heme to form the protonated heme structure **Int3** via transition state **TS3**. A proton reshuffle leads to phenol products (**Int6**) or alternatively the NO₂ radical attacks the *ipso*-position to form nitrated Trp products (**Int5**).

For the small model complex, a small barrier of $\Delta E^{\ddagger} + ZPE = 12.2$ kcal mol⁻¹ for peroxynitrite dissociation into CpdlI and NO₂ was found via ²TS1_A. Much larger barriers are

found for the next steps in the mechanism, namely for the attack of NO₂ on the substrate (via ²TS_{4A}) or the oxo attack on the substrate (via ²TS_{2A}). With the small model complex, we find pathway 1 is favored over pathway 2, whereby ²TS_{2A} is calculated to be higher in energy than ²TS_{4A} by at least 4 kcal mol⁻¹, Supporting Information (Tables S4 and S6). Subsequently, the full mechanism was calculated for nitration at the C⁴-, C⁵-, C⁶- and C⁷-positions of L-Trp using the large model **B**. Furthermore, as substrate is bound in the substrate binding pocket with different residues on both sides, we considered attacks on either side of the aromatic ring, which are designated as the *endo*- and *exo*-pathways. In addition, to these two pathways, we also tested a direct attack of heme-bound peroxynitrite onto these carbon-positions of L-Trp to form intermediate **Int4** using model **A**. However, this pathway (pathway 3; Supporting Information Figure S20) is very high in energy and did not lead to NO₂ transfer to substrate, but instead the peroxynitrite structure stayed intact. As such, peroxynitrite will need to be activated first and split into NO₂ and an iron(IV)-oxo species in order to be reactive. Consequently, pathway 3 was ruled out for further studies. We also attempted a reaction where NO₂ attacks the substrate in structure ^{2,4}Int_{2A} to form either **Int4_A** or **Int5_A** directly. However, no viable low energy pathways were found either.

Subsequently, we focused on the large model structure (model **B**), and calculated the full reaction landscape starting from a peroxynitrite bound structure. The potential energy landscape for the reaction of the iron(III)-peroxynitrite complex with L-Trp on the lowest energy doublet and quartet spin states for pathway 1 for attack on the C⁴-*exo* position are given in Figure 3. The reactant states (**Re_B**) of the heme active site with substrate bound are in a doublet spin ground state as expected for iron(III) complexes, while the quartet spin state is $\Delta G = 3.6$ kcal mol⁻¹ higher in free energy. There is a small barrier of $\Delta G^\ddagger = 3.3$ kcal mol⁻¹ via ²TS_{1B} for homolytic O–O cleavage to form iron(IV)-oxo and an NO₂[•] radical. The quartet spin barrier is much higher in energy, i.e. $\Delta G^\ddagger = 14.5$ kcal mol⁻¹, and therefore the first reaction step will take place on a dominant doublet spin state surface.

The peroxynitrite-cleaving transition state geometries have substantially elongated O–O distances of 1.902 (1.766) Å for ²TS_{1B} (⁴TS_{1B}) and visualization of the imaginary frequency shows an O–O bond cleavage mode. The first reaction step is virtually thermoneutral on the doublet spin state and brings the quartet spin state to within a fraction of a kcal mol⁻¹. This is not surprising as, electronically, ^{4,2}Int_{1B} can be characterized as an iron(IV)-oxo heme complex, i.e. Compound II (CpdII), with an NO₂ radical in close proximity. Indeed the group spin densities (Supporting Information) of ^{4,2}Int_{1B} show a full spin on the NO₂ group. The molecular valence orbitals of ^{4,2}Int_{1B} (shown in Figure S29, Supporting Information) confirm that CpdII has an electronic configuration with two unpaired electrons in π^*

orbitals (π^*_{xz} and π^*_{yz}) for the antibonding interaction of a 3d iron orbital with a 2p on the oxo group. The π^*_{yz} orbital mixes somewhat with a $\pi^*_{NO_2}$ orbital and hence there is a small but weak long-range interaction between CpdII and NO₂. The heme-based a_{2u} orbital is doubly occupied in CpdII, in contrast to CpdI where it is singly occupied.

Thus, the two unpaired electrons in π^*_{xz} and π^*_{yz} are ferromagnetically coupled to the NO₂[•] radical (in orbital $\pi^*_{NO_2}$) into an overall quartet spin state or anti-ferromagnetically coupled into an overall doublet spin state. As such, there are two different spin-configurations for the CpdII structure with overall doublet and quartet spin. These two configurations are close in energy and indeed we find them within 1 kcal mol⁻¹.

In previous work on P450 reactivity by Compound I (CpdI), i.e. the iron(IV)-heme cation radical species which is the one electron oxidized form of CpdII, it was found to also have close lying doublet and quartet spin state structures, but with $\pi^*_{xz} \uparrow \pi^*_{yz} \uparrow a_{2u} \uparrow$ configuration.^{74–76} Thus, reaction mechanisms with substrates were shown to proceed on competing spin-state surfaces with competing rate-constants. Moreover, examples have been highlighted in the past, where product distributions were different on each of the spin-state surfaces.^{77,78} Therefore, we calculated the full mechanism of L-Trp nitration on the doublet and quartet spin states for all reactions.

We attempted to optimize the geometry of the product from heterolytic cleavage of peroxynitrite to give either ^{4,2}CpdI + NO₂⁻ or an iron(III)-oxo(heme) + NO₂[•]. However, in both cases when we swapped molecular orbitals in ^{4,2}Int_{1B}, the SCF converged back to the original conformation. Therefore, the heterolytic cleavage pathways are high in energy and are unlikely scenarios for the TxtE and can be ruled out. As such we do not expect either NO₂[•] or NO₂⁻ to be involved in the substrate nitration reaction mechanism.

Subsequently, we investigated the addition of NO₂ to L-Trp substrate from the ^{2,4}Int_{1B} structures and, as an example, we show the calculated mechanism for activation of the C⁴-*exo*-position of L-Trp by CpdII in Figure 3. The C–N bond formation transition states take place via barriers ^{2,4}TS_{4B,C4-exo}. On both the doublet and quartet spin state surfaces, small barriers that are within $\Delta G^\ddagger = 1$ kcal mol⁻¹ of ^{4,2}Int_{1B} are found. Nevertheless, both transition states were characterized as shown in Figure 3 with a single imaginary frequency of i178/i179 cm⁻¹ representing a C–NO₂ vibrational stretch. As the transition states represent NO₂ transfer to L-Trp and no electron is transferred to CpdII, there are little changes to the geometry of the CpdII part as compared between ^{2,4}Int_{1B} and ^{2,4}TS_{4B,C4-exo}, and geometrically those parts of the structure are the same.

Table 1. Relative energies of NO₂ transfer to various positions of the L-Trp substrate.^a

Pathway	² TS _{4B} (⁴ TS _{4B})	² Int _{4B} (⁴ Int _{4B})	² TS _{5B} (⁴ TS _{5B})	² Int _{5B} (⁴ Int _{5B})
C ⁴ -exo	>0.0 (>0.0)	-3.5 (-3.5)	>-3.0 (>-3.0)	-58.0 (-23.2)
C ⁴ -endo	6.6 (>0.0)	6.4 (-1.4)	11.0 (15.0)	-49.2 (-14.9)
C ⁵ -exo	0.9 (0.9)	-0.6 (0.0)	16.9 (17.9)	-49.6 (-15.8)
C ⁵ -endo	>3.0 (>0.0)	2.7 (-4.2)	8.3 (10.3)	-57.7 (-23.6)
C ⁶ -exo	0.6 (>0.0)	0.3 (-0.4)	6.1 (6.6)	-40.8 (-46.9)
C ⁶ -endo	3.2 (2.2)	0.2 (1.1)	6.4 (6.1)	-45.9 (-40.8)
C ⁷ -exo	2.4 (2.4)	0.2 (0.2)	2.7 (4.5)	-60.9 (-56.1)
C ⁷ -endo	>0.0 (>0.0)	-1.6 (-1.5)	1.8 (3.8)	-56.9 (-27.2)

^a Values in kcal mol⁻¹ represent ΔE+ZPE data with energies obtained at UB3LYP/BS1//UB3LYP/BS2 relative to ²Re_B.

The transition state is a pure NO₂[•] attack on the aromatic ring of L-Trp and no electron transfer to the heme takes place. The shallow intermediate (^{4,2}Int_{4B,C⁴-exo}) is followed by another low-energy barrier, namely the hydrogen atom transfer from L-Trp to CpdII, which could not be characterized for the C⁴-exo pathway and hence will proceed very fast. As such, these reactions will be fast and efficient, and it is unlikely that NO₂[•] will drift out of the substrate binding pocket if substrate is present. Obviously, it is important that NO₂[•] does not drift out of the protein pocket as it is known to activate aromatic amino acids and could react with enzymatic residues if the substrate is not positioned appropriately.³⁵ In particular, the substrate binding pocket of the 4TPO pdb file²⁴ is lined with several aromatic residues, e.g. Phe₇₉, Trp₈₂, Tyr₈₉, Trp₂₉₇ and Phe₃₉₅. Clearly, peroxynitrite activation and the production of NO₂ should not lead to nitration of any of these active site residues. Therefore, NO₂ should be used by the protein as soon as it is formed and, based on the low barriers found here, they clearly implicate a rapid substrate activation.


As our proposed mechanism of L-Trp nitration by P450 enzymes suggests the formation of a free NO₂ radical after homolytic cleavage of peroxynitrite, we decided to explore pathways leading to C⁴-, C⁵-, C⁶- and C⁷-nitration for model **B** and the results are summarized in Table 1. Of course, these mechanisms have two stereochemical isomers, which are designated as the *exo*- and *endo*-pathways. Optimized geometries of all ^{2,4}TS_{4B} transition states are given in the Supporting Information Figure S25, while the one for C⁴-exo is given above in Figure 3. As this reaction step proceeds away from the heme center and no electron transfer takes place, it means that the heme CpdII distances, i.e. the Fe–O and Fe–S interactions, are very close for all structures. A little more variation is seen in the C–N bond that is being formed in the transition state. The shortest distances are found for C⁵-activation whereby the *endo*-pathway gives the shortest distances of 1.714 Å, while those for the *exo*-pathway are 1.721 Å. By contrast, much longer C–N distances are found for the C⁷-activation with values of 1.830 / 1.831 Å for ²TS_{4B,C⁷-endo} / ⁴TS_{4B,C⁷-endo} and 1.827 / 1.842 Å for ²TS_{4B,C⁷-exo} / ⁴TS_{4B,C⁷-exo}.

Next, we looked at the energetics for NO₂ transfer to the C⁴-, C⁵-, C⁶- and C⁷-positions of the L-Trp substrate

through attack from the *exo* and *endo* directions. We located C–N bond formation transition states (^{2,4}TS_{4B}) and the subsequent local minima (^{2,4}Int_{4B}), as well as the *ipso*-hydrogen atom abstraction transition state (^{2,4}TS_{5B}) by CpdII leading to the nitrated product complexes. ^{2,4}Int_{5B} and the energies with respect to ²Re_B for all pathways are given in Table 1. Several barriers for C–N bond formation via TS_{4B} are very low in energy and could not be located precisely, but geometry scans find them to be less than 1 kcal mol⁻¹ above the Int_{4B} intermediate. These pathways should lead to addition of NO₂ to L-Trp at the C⁴-exo, C⁴-endo, C⁵-endo, C⁶-exo and C⁷-endo positions. As a matter of fact, most TS_{4B} barriers are found within a small window of 4 kcal mol⁻¹ and only doublet spin C⁴-endo is higher in energy, i.e. its transition state is about 7 kcal mol⁻¹ above ²Int_{4B}.

The NO₂ addition local minimum (^{2,4}Int_{4B}) is followed by hydrogen abstraction from the *ipso*-position by CpdII to form nitrated Trp and ferric-hydroxide. Due to the position of the substrate in the binding pocket, in several cases substantial hydrogen abstraction barriers are found. Thus, the C⁶- and C⁷-positions of L-Trp are pointing toward the iron(IV)-oxo group of the heme and hence, when NO₂ attacks the substrate at these positions, their hydrogen atoms can easily be abstracted by CpdII. Indeed, TS_{5B} barriers for C⁶- and C⁷-nitration barriers of less than 7 kcal mol⁻¹, with respect to isolated reactants, are those mostly found. By contrast, the C⁵-exo position points away from the heme and consequently hydrogen atom transfer is difficult and high barriers are obtained as a result of major structural rearrangements.

Finally, we investigated the possible aromatic hydroxylation reaction of P450 TxtE (bottom-right pathway in Scheme 4). As shown before, the aromatic hydroxylation by a metal(IV)-oxo species starts with an electrophilic addition of the oxo to the aromatic ring, followed by proton shuttle from the aromatic *ipso*-position to a nearby proton acceptor group, such as the nitrogen atoms of the porphyrin ring, prior to relay of the proton to the oxygen atom to form hydroxylated product.^{79–81}



Scheme 5. Comparative catalytic cycles of P450 nitration and P450 mono-oxygenases.

This mechanism, i.e. pathway 2 in Scheme 4, was found to be high in energy for the quartet spin state ($\Delta E^\ddagger + \text{ZPE} > 30 \text{ kcal mol}^{-1}$ for ${}^4\text{TS}_{2\text{A}}$ and ${}^4\text{TS}_{3\text{A}}$) and, although somewhat lower on the doublet spin state surface ($>25 \text{ kcal mol}^{-1}$), is still much higher in energy than the NO_2^\bullet transfer step to the substrate. Therefore, substrate hydroxylation will not be a feasible pathway in the presence of free NO_2^\bullet radicals. This is in agreement with experimental work on P450 TxtE that failed to identify hydroxylated L-Trp products.

An alternative to the electrophilic aromatic hydroxylation, that started with a hydrogen atom abstraction from the substrate by the iron(IV)-oxo species in **Int2**, was also investigated. In particular, the hydrogen atom abstraction from the C⁶ position was explored for the large model system. However, high energy hydrogen atom abstraction free energies of activation of $\Delta G^\ddagger = 23.5 \text{ kcal mol}^{-1}$ and $22.8 \text{ kcal mol}^{-1}$ for doublet and quartet, respectively, were obtained (Supporting Information Figure S27). Therefore, a direct hydrogen atom abstraction from the aromatic ring of L-Trp can be ruled out as a viable mechanism as expected given the high C–H bond strength.

Overall, the work shows that the aromatic nitration by P450 TxtE is triggered by a reaction of NO with an iron(III)-superoxo species to form iron(III)-peroxynitrite that releases NO_2^\bullet rapidly. These NO_2^\bullet radicals react with the indole ring of L-Trp through addition and the nearby CpdII species abstracts the *ipso*-hydrogen atom from the 4-nitrotryptophan products. However, as a free NO_2^\bullet is formed, this means regioselectivity is determined by the tightness of substrate binding and the orientation of Trp in

the substrate binding pocket. How this selectivity is determined, we discuss in further detail below.

DISCUSSION.

In this work a combined experimental and computational study on the P450 aromatic nitration enzyme TxtE is reported. These studies focus on the catalytic reaction mechanism and the nature of the selectivity of the reaction. Overall, our studies show that the mechanism is distinctly different from conventional P450 mono-oxygenases, where an active species (Compound I, CpdI) is initially formed and subsequently reacts with substrates.^{1–5} To explain the differences in the catalytic reaction cycle of P450 TxtE and P450 mono-oxygenases in general, we compare the two reaction mechanisms side-by-side in Scheme 5. Both catalytic cycles start from an iron(III)-heme complex that is linked to the protein through a thiolate bridge with a cysteine residue. The sixth coordination site of the iron(III) is occupied by a water molecule in the resting state (structure **A** in Scheme 5). Upon substrate binding the water molecule is released and the metal complex undergoes a spin-state change from low-spin to high-spin. DFT calculations showed that the high-spin is preferential over the low-spin in the 5-coordinate iron(III) complex, whereas the spin-state ordering is reversed in 6-coordinate environments.⁸² Thereafter, the complex is reduced by the redox partner and binds molecular oxygen (complex **B** in Scheme 5). Beyond complex **B**, the catalytic cycles of P450 mono-oxygenases and P450 nitration enzymes diverge.




Figure 4. Extracts of the crystal structure coordinates of P450 TxtE (left) and P450cam (right) as taken from the 4TPO and 3WRH pdb files. Amino acids are labeled as in the pdb files.

Thus, the mono-oxygenases reduce structure **B** and after two proton transfer steps generate a high-valent iron(IV)-oxo heme cation radical species called CpdI. The P450 TxtE isozymes, by contrast, bind a molecule of NO that reacts with the iron(III)-superoxo intermediate (**B**) to form an iron(III)-peroxynitrite intermediate.

As discussed above, several pathways for nitration of L-Trp substrate were calculated here and our lowest energy mechanism involves a reaction that starts with the iron(III)-peroxynitrite splitting into Compound II (CpdII), i.e. iron(IV)-oxo heme and NO_2^\bullet radical. The latter then attacks the substrate at the C⁴-position and the *ipso*-hydrogen atom at C⁴ moves to CpdII to generate an iron(III)-hydroxo complex. A proton transfer from the protein or solvent then brings the catalytic cycle back into the resting state. At this stage it is not clear whether this proton transfer step happens prior to, or after, product release.

The catalytic cycles of P450 nitration and mono-oxygenation in Scheme 5 show dramatic differences. Thus, P450 TxtE would require only a single electron from a redox partner and no proton transfer to generate an iron(III)-peroxynitrite as the active species, whereas P450 mono-oxygenases need two electrons and two external protons to form CpdI. However, one proton transfer is needed in P450 TxtE to convert protonated CpdII back to the resting state. To understand these differences in proton relay mechanisms during the catalytic cycle of P450 TxtE and general P450 mono-oxygenases, we analyzed structures of representative nitration and mono-oxygenase enzymes (Figure 4). In particular, Figure 4 displays an extract of the active site structures of P450 TxtE (4TPO pdb as a representative of a nitrating P450)²⁴ and P450 CAM (3WRH pdb as a mono-oxygenase P450 structure).^{83,84}

P450 CAM hydroxylates camphor regio- and stereoselectively at the C⁴-position and its mechanism has been extensively studied with experimental and computational

methods.^{85–88} As rationalized and discussed previously,^{73,87} P450 CAM has a number of active site residues involved in proton-relay during the catalytic cycle. Firstly, a buried Glu₃₆₆ residue is connected via a chain of water molecules to the alcohol group of Thr₂₅₂ that is located just above the heme. The Glu₃₆₆ residue is believed to deliver the first proton in the catalytic cycle of P450 mono-oxygenases,⁸⁹ while Asp₂₅₁ delivers the second one. The Asp₂₅₁ residue points out of the active site, but computational modelling showed that a rotation of the side chain brings the carboxylate group close to the Thr₂₅₁ alcohol group, and thus Asp₂₅₁ can deliver a second proton into the substrate binding pocket to complete the formation of CpdI.

An analysis of the TxtE structure of 4TPO shows that the Glu₃₆₆ residue is missing and is replaced by a Gln residue. Nevertheless, the Gln environment is similar to that of Glu₃₆₆ in P450 CAM, namely at the end of a cavity surrounded by apolar residues, such as Leu and Ile. Interestingly, Val₂₅₃ is not seen in P450 TxtE and is replaced by another Thr residue (Thr₂₅₁). Nevertheless, it is clear from the TxtE structure that one proton source is lacking in the active site. Therefore, proton transfer in P450 TxtE will be considerably slower than in P450 CAM, where one proton may already be stored on the Glu₃₆₆ residue prior to the start of the catalytic cycle, and which should lead to efficient protonation of the iron(II)-peroxo to form an iron(III)-hydroperoxo complex.

In P450 CAM the second proton in the catalytic cycle is proposed to be delivered through the rotation of Asp₂₅₁. In the TxtE structure, however, this Asp residue is missing and a proline residue takes its position.




Figure 5. Valence bond scheme for the electron migrations during the conversion of the ferric-superoxo and NO into CpdII and NO₂ via a peroxynitrite intermediate.

Therefore, the proton transfer residues in P450 TxtE have been mutated with respect to typical mono-oxygenase structures, and consequently proton transfer into the active site for the conversion of the iron(III)-superoxo species into CpdI is hampered. This should give the iron(III)-superoxo species a relatively long lifetime until NO is delivered and can form a heme-peroxynitrite complex.

The electron transfer processes that happen in the reaction of the iron(III)-superoxo species with NO to form NO₂ via a peroxynitrite intermediate are given in Figure 5 as a valence bond (VB) diagram. VB schemes have been extensively used to explain electron transfer processes in reaction mechanisms and, in particular, were found to be useful to explain regio- and chemoselectivities of reaction processes and make predictions on what bonds are most likely activated in a substrate.⁹⁰⁻⁹³ These VB structures give a bond orbital occupied by two electrons as a line separated by two dots, while individual (unpaired) electrons are given as dots. Thus, the isolated iron(III)-superoxo structure has an open-shell singlet ground state with orbital occupation $\delta_{xz-yz}^2 3d_{xz}^2 \pi_{yz}^2 \pi_{yz}^* \pi_{OO,yz}^2 \pi_{OO,yz}^* \pi_{OO,xz}^2 \pi_{OO,xz}^*$.⁷⁶ The π_{OO} and π_{OO}^* orbitals represent the bonding and antibonding orbitals along the superoxo bond in the xz and yz planes. The metal-type orbitals contain the doubly occupied δ_{xz-yz} orbital, which is a nonbonding orbital in the plane of the heme with lobes pointing toward the meso-carbon atoms. Then there are three electrons along the Fe-O bond in $\pi_{yz}(2)$ and $\pi_{yz}^*(1)$ for the bonding and antibonding interaction of the metal $3d_{yz}$ with a $2p_y$ on oxygen. This interaction is described as a two-center-three-electron bond between the Fe and O atoms in Figure 5. In the xz plane the $3d_{xz}$ iron orbital is not involved in interactions with any of the oxygen atoms as the peroxo group is in that plane so, in contrast to CpdII, it is a doubly occupied atomic orbital in the iron(III)-superoxo form: $3d_{xz}^2$. The approaching NO molecule has one unpaired electron on the nitrogen atom. Upon attack of the NO on the superoxo radical an N-O bond is formed by the pairing up of the $\pi_{OO,xz}^*$ and π_N radicals to form the peroxynitrite π_{NO} bond with two electrons. At the same time, the two electrons from the $\pi_{OO,yz}$ orbital in the iron(III)-superoxo structure become the lone-pair electrons on bridging oxygen of the peroxynitrite group: lp_O . This then generates the iron(III)-

peroxynitrite intermediate, i.e. the reactant complex **Re** in Figure 3 and Scheme 4, with orbital configuration $\delta_{xz-yz}^2 3d_{xz}^2 \pi_{yz}^2 \pi_{yz}^* \pi_{OO,xz}^2 \pi_{OO,xz}^* \pi_{NO}^2 lp_O^2$. Consequently, the bonding and antibonding type electrons involved in the superoxo bond in the iron(III)-superoxo species are redistributed over the chemical system and a single bond remains between the two oxygen atoms, i.e. dioxygen configuration $\pi_{OO,xz}^2 \pi_{OO,xz}^*$.

Next, the O-O bond breaks homolytically to form NO₂ and a CpdII fragment by moving one electron from the $\pi_{OO,xz}$ orbital in **Re** into the Fe-O bond. This electron pairs up with the two electrons of the $3d_{xz}$ orbital on the iron to form the new two-center three-electron bond with $\pi_{xz}^2 \pi_{xz}^*$ configuration. CpdII is in a triplet spin ground state,⁹¹ with $\pi_{xz}^* \pi_{yz}^*$ configuration that is either antiferromagnetically coupled to a radical on NO₂ in a doublet spin state or ferromagnetically coupled into a quartet spin state. Indeed, ^{4,2}**Inti** are close in energy, while in ^{4,2}**Re** the doublet and quartet spin states are well separated. The calculations reported in this work show that a reaction of iron(III)-superoxo with NO rapidly forms iron(III)-peroxynitrite complexes. Moreover, the subsequent reaction for homolytic cleavage of peroxynitrite to form NO₂ and CpdII has a calculated barrier of only $\Delta G^\ddagger = 3.3$ kcal mol⁻¹ and hence should proceed rapidly. These results are consistent with recent work of the Moënne-Loccoz group on a biomimetic non-heme Cu(II)-peroxynitrite complex that was found to homolytically cleave into NO₂ and a copper(II)-oxo species.⁷²⁻⁹⁴

As shown in this work, homolytic cleavage of peroxynitrite gives CpdII and free NO₂, which can add to the π -system of the substrate. These radicals can attack and react with the substrate. To find out if a bare indole group in the gas-phase would have a preference for nitration of a specific carbon site in L-Trp, we calculated the C-N bond formation energy by addition of NO₂[•] to methylindole, see Figure 6. As can be seen in all cases, the addition of NO₂[•] to methylindole is endothermic by 10.5 – 15.3 kcal mol⁻¹ in the gas-phase with a preference for addition at the C⁴-position. As such, if all carbon atoms of L-Trp are accessible in the protein then a reaction with NO₂ should give a dominant C⁴-nitration.




Figure 6. (a) UB3LYP/6-31G* calculated C–N bond formation energies in a bare methylindole molecule. Values are $\Delta E + \text{ZPE}$ data in kcal mol⁻¹. (b) Electric field effect (with E_{field} in units of 0.001 au) on the relative energies of NO_2 addition to methylindazole along the z-axis.

Indeed experimental studies on L-Trp activation by P450 TxtE have confirmed that this is the only detectable product.^{22–24,32} The results of the calculations reported in Table 1 above show that the lowest NO_2 addition barrier is for the C⁴-exo pathway. However, the table shows several alternative pathways to C⁵-, C⁶- and C⁷-nitration to be close in energy. Hence, a mixture of products is expected in an open substrate binding pocket. Consequently, a tight binding of substrate and oxidant that brings NO_2^* and the target position of substrate close is essential for P450 TxtE.

Interestingly, the addition of NO_2 to the aromatic ring of methylindole is endothermic by well over 10 kcal mol⁻¹, whereas the same step in the large enzymatic structure is almost thermoneutral as seen by the energy differences for model **B** of **Int1** and **Int4** in Figure 3. By contrast, for the small model complex without the protein environment (model **A**) an energy of $\Delta E + \text{ZPE} = 19.4$ (doublet) or 18.6 (quartet) kcal mol⁻¹ is obtained for the same reaction step. Therefore, the protein environment stabilizes complex **Int4** dramatically and enables a favorable substrate nitration reaction with low barriers.

To find out how the long-range interactions stabilize the NO_2 adduct **Int4** we took the methylindole structures and

calculated their energies through the addition of an external electric field.^{95,96} Electric fields were applied to the molecular x-, y-, and z-axes, whereby the y-axis is along the C– NO_2 bond that is being formed and the z-axis along the C–*ipso*-H coordinate as shown on the top-right in Figure 6. These applied electric fields are of a magnitude similar to those used in vibrational Stark effect spectroscopy measurements on enzymatic systems and hence mimic realistic perturbations.⁹⁷ With a field along the x- and y-axis small energy changes are seen, but along the z-axis these changes are major (see Figure 6b). Therefore, the barrier height of the reaction is linked with the movement of the C–H bond from in-plane with respect to the Trp group to make space for the in-coming nitro group. Thus, an applied electrical field along the z-axis stabilizes the NO_2 adduct with methylindole dramatically and changes the reaction energy for adduct formation from endothermic by 10 kcal mol⁻¹ to exothermic by 10 kcal mol⁻¹. Particularly, the C⁴-addition pathway is stabilized with an electric field effect along the z-axis and enhances the selectivity of the reaction. As a consequence, the protein is able to stabilize and drive an otherwise unfavorable reaction through an induced polarized environment. The substrate binding pocket has many

polar residues, e.g. Arg₅₉ and Glu₃₉₄ that bind the carboxylate and ammonium groups of the substrate.²⁴ In addition, the side chains of the polar residues Met₈₈, Tyr₈₉, Met₁₇₃ and Asn₂₉₃ contact the indole group of the Trp substrate and through electrostatic and induced dipole moment interactions may affect the selectivity of the reaction, but this will need to be confirmed by future mutation studies.

CONCLUSIONS.

In this work a combined experimental and computational study is presented for the possible reaction mechanisms of aromatic nitration by a P450 isozyme. We show that dioxygen binds first to the heme iron, followed by NO to generate an iron(III)-peroxynitrite species. We set up several active site model complexes with bound L-Trp and investigated substrate activation pathways leading to nitration as well as to oxidation of the substrate. We found low-energy barriers for the homolytic cleavage of iron(III)-peroxynitrite into NO₂ and CpdII. Consequently, the reaction of NO and O₂ on an iron(III)-heme center is expected to generate free NO₂ radicals efficiently. We then explored NO₂ attack on L-Trp and found pathways for activation at the C⁴, C⁵, C⁶ and C⁷-positions. Although NO₂ addition is generally a low-energy process, many pathways encounter relatively high energy hydrogen atom transfer steps to form products. As such, substrate positioning and binding must be essential in P450 TxtE as indeed is observed experimentally,³³ where a point mutation in a dynamic loop forming the active site lid caused a complete switch of nitration regiochemistry from C⁴ to C⁵.

ASSOCIATED CONTENT

Supporting Information. Detailed experimental data, including experimental set-up of double mixing experiments, spectroscopic and mass spectrometric data. In addition, full computational data with all optimized geometries on model A and B for the substrate hydroxylation and nitration of L-Trp by an iron(III)-peroxynitrite structure is given in the Supporting Information. The data contain absolute and relative energies, group spin densities and charges, and Cartesian coordinates of all optimized geometries. In addition, constraint geometry scan, drawings of optimized geometries and intrinsic reaction coordinate scans are given. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

* sam.devisser@manchester.ac.uk (SPdV), G.L.Challis@warwick.ac.uk (GLC), Nigel.Scrutton@manchester.ac.uk (NSS).

Present Addresses

†Department of Chemistry, Faculty of Natural & Mathematical Sciences, King's College London, Britannia House, 7 Trinity Street, London SE1 1DB, United Kingdom.

Author Contributions

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript.

‡ These authors contributed equally.

Funding Sources

The Biotechnology and Biological Sciences Research Council (BBSRC) is thanked for support to NSS, AWM and GLC under grant code BB/H006265/1 and BB/H006281/1.

ACKNOWLEDGMENT

MQEM thanks the Malaysian Government and Islamic Science University of Malaysia (USIM) for a studentship. The Punjab Education Endowment Fund (PEEF) in Pakistan is acknowledged for a PhD scholarship to HAS.

ABBREVIATIONS

BDE, bond dissociation energy; CpdI, Compound I; CpdII, Compound II; P450, cytochrome P450; DFT, density functional theory; DTT, dithiothreitol; HP, high performance; VB, valence bond; ZPE, zero-point energy.

REFERENCES

- (1) Sono, M.; Roach, M. P.; Coulter, E. D.; Dawson, J. H. Heme-containing oxygenases. *Chem. Rev.* **1996**, *96*, 2841–2888.
- (2) Ortiz de Montellano, P. R., Ed., *Cytochrome P450: Structure, Mechanism and Biochemistry*, 3rd Ed., Kluwer Academic/Plenum Publishers, New York, **2005**.
- (3) Kadish, K. M.; Smith, K. M.; Guillard, R., Eds., *Handbook of Porphyrin Science*, World Scientific Publishing Co., New Jersey, **2010**.
- (4) Munro, A. W.; Girvan, H. M.; McLean, K. J. Variations on a (t)heme—novel mechanisms, redox partners and catalytic functions in the cytochrome P450 superfamily *Nat. Prod. Rep.* **2007**, *24*, 585–609.
- (5) Ortiz de Montellano, P. R. Hydrocarbon hydroxylation by cytochrome P450 enzymes. *Chem. Rev.* **2010**, *110*, 932–948.
- (6) Krauser, J. A.; Guengerich, F. P. Cytochrome P450 3A4-catalyzed testosterone 6β-hydroxylation stereochemistry, kinetic deuterium isotope effects, and rate-limiting steps. *J. Biol. Chem.* **2005**, *280*, 19496–19506.
- (7) Spinello, A.; Pavlin, M.; Casalino, L.; Magistrato, A. A dehydrogenase dual hydrogen abstraction mechanism promotes estrogen biosynthesis: can we expand the functional annotation of the aromatase enzyme? *Chem. Eur. J.* **2018**, *24*, 10840–10849.
- (8) Greule, A.; Stok, J. E.; De Voss, J. J.; Cryle, M. J. Unrivalled diversity: the many roles and reactions of bacterial cytochromes P450 in secondary metabolism. *Nat. Prod. Rep.* **2018**, *35*, 757–791.
- (9) Fessner, N. D. P450 monooxygenases enable rapid late-stage diversification of natural products via C–H bond activation. *ChemCatChem*. **2019**, *11*, 2226–2242.
- (10) Spinello, A.; Ritacco, I.; Magistrato, A. The catalytic mechanism of steroidogenic cytochromes P450 from all-atom simulations: Entwinement with membrane environment, redox partners, and post-transcriptional regulation. *Catalysts* **2019**, *9*, 81.
- (11) Grogan, G. Cytochromes P450: exploiting diversity and enabling application as biocatalysts. *Curr. Opin. Chem. Biol.* **2011**, *15*, 241–248.

- (12) Wei, Y.; Ang, E. L.; Zhao, H. Recent developments in the application of P450 based biocatalysts. *Curr. Opin. Chem. Biol.* **2018**, *43*, 1-7.
- (13) Li, Z.; Jiang, Y.; Guengerich, F. P.; Ma, L.; Li, S.; Zhang, X. W. Engineering cytochrome P450 enzyme systems for biomedical and biotechnological applications. *J. Biol. Chem.* **2020**, *295*, 833-849.
- (14) Poulos, T. L. Heme enzyme structure and function. *Chem. Rev.* **2014**, *114*, 3919-3962.
- (15) Sevrioukova, I. F.; Poulos, T. L. Understanding the mechanism of cytochrome CYP450 3A4: Recent advances and remaining problems. *Dalton Trans.* **2013**, *42*, 3116-3126.
- (16) Green, M. T. C-H bond activation in heme proteins: the role of thiolate ligation in cytochrome P450. *Curr. Opin. Chem. Biol.* **2009**, *13*, 84-88.
- (17) Meunier, B.; de Visser, S. P.; Shaik, S. Mechanism of oxidation reactions catalyzed by cytochrome P450 enzymes. *Chem. Rev.* **2004**, *104*, 3947-3980.
- (18) Denisov, I. G.; Makris, T. M.; Sligar, S. G.; Schlichting, I. Structure and chemistry of cytochrome P450. *Chem. Rev.* **2005**, *105*, 2253-2277.
- (19) Huang, X.; Groves, J. T. Oxygen activation and radical transformations in heme proteins and metalloporphyrins. *Chem. Rev.* **2018**, *118*, 2491-2553.
- (20) Dunbar, K. L.; Scharf, D. H.; Litomska, A.; Hertweck, C. Enzymatic carbon-sulfur bond formation in natural product biosynthesis. *Chem. Rev.* **2017**, *117*, 5521-5577.
- (21) Guengerich, F. P.; Yoshimoto, F. K. Formation and cleavage of C-C bonds by enzymatic oxidation-reduction reactions. *Chem. Rev.* **2018**, *118*, 6573-6655.
- (22) Barry, S. M.; Kers, J. A.; Johnson, E. G.; Song, L.; Aston, P. R.; Patel, B.; Krasnoff, S. B.; Crane, B. R.; Gibson, D. M.; Loria, R.; Challis, G. L. Cytochrome P450-catalyzed L-tryptophan nitration in thaxtomin phytotoxin biosynthesis. *Nat. Chem. Biol.* **2012**, *8*, 814-816.
- (23) Yu, F.; Li, M.; Xu, C.; Wang, Z.; Zhou, H.; Yang, M.; Chen, Y.; Tang, L.; He, J. Structural insights into the mechanism for recognizing substrate of the cytochrome P450 enzyme TxtE. *PLoS One* **2013**, *8*, e81526.
- (24) Dodani, S. C.; Cahn, J. K. B.; Heinisch, T.; Brinkmann-Chen, S.; McIntosh, J. A.; Arnold, F. H. Structural, functional, and spectroscopic characterization of the substrate scope of the novel nitrating cytochrome P450 TxtE. *ChemBioChem.* **2014**, *15*, 2259-2267.
- (25) Girvan, H. M.; Munro, A. W. Applications of microbial cytochrome P450 enzymes in biotechnology and synthetic biology. *Curr. Opin. Chem. Biol.* **2016**, *31*, 136-145.
- (26) Dodani, S. C.; Kiss, G.; Cahn, J. K. B.; Su, Y.; Pande, V. S.; Arnold, F. H. Discovery of a regioselectivity switch in nitrating P450s guided by molecular dynamics simulations and Markov models. *Nat. Chem.* **2016**, *8*, 419-425.
- (27) Zuo, R.; Zhang, Y.; Jiang, C.; Hackett, J. C.; Loria, R.; Bruner, S. D.; Ding, Y. Engineered P450 biocatalysts show improved activity and regio-promiscuity in aromatic nitration. *Sci. Rep.* **2017**, *7*, 842.
- (28) Tomita, H.; Katsuyama, Y.; Minami, H.; Ohnishi, Y. Identification and characterization of a bacterial cytochrome P450 monooxygenase catalyzing the 3-nitration of tyrosine in rifomycin biosynthesis. *J. Biol. Chem.* **2017**, *292*, 15859-15869.
- (29) Jiang, G.; Zuo, R.; Zhang, Y.; Powell, M. M.; Zhang, P.; Hylton, S. M.; Loria, R.; Ding, Y. One-pot biocombinatorial synthesis of herbicidal thaxtomins. *ACS Catal.* **2018**, *8*, 10761-10768.
- (30) Zuo, R.; Ding, Y. Direct aromatic nitration system for synthesis of nitrotryptophans in *Escherichia coli*. *ACS Synth. Biol.* **2019**, *8*, 857-865.
- (31) Buddha, M. R.; Tao, T.; Parry, R. J.; Crane, B. R. Regioselective nitration of tryptophan by a complex between bacterial nitric-oxide synthase and tryptophanyl tRNA synthetase. *J. Biol. Chem.* **2004**, *279*, 49567-49570.
- (32) Greule, A.; Stok, J. E.; De Voss, J. J.; Cryle, M. J. Unrivalled diversity: the many roles and reactions of bacterial cytochromes P450 in secondary metabolism. *Nat. Prod. Rep.* **2018**, *35*, 757-791.
- (33) Alkhalaf, L. M.; Barry, S. M.; Rea, D.; Gallo, A.; Griffiths, D.; Lewandowski, J. R.; Fulop, V.; Challis, G. L. Binding of distinct substrate conformations enables hydroxylation of remote sites in Thaxtomin D by cytochrome P450 TxtC. *J. Am. Chem. Soc.* **2019**, *141*, 216-222.
- (34) Schopfer, F. J.; Baker, P. R. S.; Freeman, B. A. NO-dependent protein nitration: a cell signaling event or an oxidative inflammatory response? *Trends Biochem. Sci.* **2003**, *28*, 646-654.
- (35) Lin, H.-L.; Kenaan, C.; Zhang, H.; Hollenberg, P. F. Reaction of human cytochrome P450 3A4 with peroxynitrite: Nitrotyrosine formation on the proximal side impairs its interaction with NADPH-cytochrome P450 reductase. *Chem. Res. Toxicol.* **2012**, *25*, 2642-2653.
- (36) Ford, P. C.; Lorković, I. M. Mechanistic aspects of the reactions of nitric oxide with transition-metal complexes. *Chem. Rev.* **2002**, *102*, 993-1017.
- (37) Herold, S.; Matsui, T.; Watanabe, Y. Peroxynitrite isomerization catalyzed by His64 myoglobin mutants. *J. Am. Chem. Soc.* **2001**, *123*, 4085-4086.
- (38) Hersleth, H.-P.; Ryde, U.; Rydberg, P.; Görbitz, C. H.; Andersson, K. K. Structures of the high-valent metal-ion haem-oxygen intermediates in peroxidases, oxygenases and catalases. *J. Inorg. Biochem.* **2006**, *100*, 460-476.
- (39) Groenhof, A. R.; Ehlers, A. W.; Lammertsma, K. Proton assisted oxygen-oxygen bond splitting in cytochrome P450. *J. Am. Chem. Soc.* **2007**, *129*, 6204-6209.
- (40) Guengerich, F. P.; Munro, A. W. Unusual cytochrome P450 enzymes and reactions. *J. Biol. Chem.* **2013**, *288*, 17065-17073.
- (41) Erdogan, H. One small step for cytochrome P450 in its catalytic cycle, one giant leap for enzymology. *J. Porph. Phthal.* **2019**, *23*, 358-366.
- (42) Dubey, K. D.; Shaik, S. Cytochrome P450: The wonderful nanomachine revealed through dynamic simulations of the catalytic cycle. *Acc. Chem. Res.* **2019**, *52*, 389-399.
- (43) Rittle, J.; Green, M. T. Cytochrome P450 compound I: capture, characterization, and C-H bond activation kinetics. *Science* **2010**, *330*, 933-937.
- (44) Lang, J.; Maréchal, A.; Couture, M.; Santolini, J. Reaction intermediates and molecular mechanism of peroxynitrite activation by NO synthases. *Biophys. J.* **2016**, *111*, 2099-2109.
- (45) Toader, V.; Xu, X.; Nicolescu, A.; Yu, L.; Bolton, J. L.; Thatcher, G. R. J. Nitrosation, nitration, and autooxidation of the selective estrogen receptor modulator raloxifene by nitric oxide, peroxynitrite, and reactive nitrogen/oxygen species. *Chem. Res. Toxicol.* **2003**, *16*, 1264-1276.
- (46) Behan, R. K.; Hoffart, L. M.; Stone, K. L.; Krebs, C.; Green, M. T. Reaction of cytochrome P450BM3 and peroxynitrite yields nitrosyl complex. *J. Am. Chem. Soc.* **2007**, *129*, 5855-5859.
- (47) Su, J.; Groves, J. T. Direct detection of the oxygen rebound intermediates, ferryl Mb and NO₂, in the reaction of metmyoglobin with peroxynitrite. *J. Am. Chem. Soc.* **2009**, *131*, 12979-12988.

- (48) Hernández-Ortega, A.; Quesne, M. G.; Bui, S.; Heyes, D. J.; Steiner, R. A.; Scrutton, N. S.; de Visser, S. P. Catalytic mechanism of cofactor-free dioxygenases and how they circumvent spin-forbidden oxygenation of their substrates. *J. Am. Chem. Soc.* **2015**, *137*, 7474–7487.
- (49) Lim, M. D.; Lorković, I. M.; Ford, P. C. The preparation of anaerobic nitric oxide solutions for the study of heme model systems in aqueous and nonaqueous media: some consequences of NO x impurities. *Methods Enzymol.* **2005**, *396*, 3–17.
- (50) Aga, R. G.; Hughes, M. N. The preparation and purification of NO gas and the use of NO releasers: the application of NO donors and other agents of nitrosative stress in biological systems. *Methods Enzymol.* **2008**, *436*, 35–48.
- (51) Cole, L. J.; Huston, W. M.; Moir, J. W. B. Delivery of nitric oxide for analysis of the function of cytochrome c'. *Methods Enzymol.* **2008**, *436*, 21–33.
- (52) Hartung, J. Organic radical reactions associated with nitrogen monoxide. *Chem. Rev.* **2009**, *109*, 4500–4517.
- (53) Quesne, M. G.; Borowski, T.; de Visser, S. P. Quantum mechanics/molecular mechanics modelling of enzymatic processes: Caveats and breakthroughs. *Chem. Eur. J.* **2016**, *22*, 2562–2581.
- (54) Ghafoor, S.; Mansha, A.; de Visser, S. P. Selective hydrogen atom abstraction from dihydroflavonol by a nonheme iron center is the key step in the enzymatic flavonol synthesis and avoids byproducts. *J. Am. Chem. Soc.* **2019**, *141*, 20278–20292.
- (55) Gaussian 09, Revision D.01, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., C. T. Wallingford, 2009.
- (56) Cantú Reinhard, F. G.; Sainna, M. A.; Upadhyay, P.; Balan, G. A.; Kumar, D.; Fornarini, S.; Crestoni, M. E.; de Visser, S. P. A systematic account on aromatic hydroxylation by a cytochrome P450 model Compound I: A low-pressure mass spectrometry and computational study. *Chem. Eur. J.* **2016**, *22*, 18608–18619.
- (57) Yang, T.; Quesne, M. G.; Neu, H. M.; Cantú Reinhard, F. G.; Goldberg, D. P.; de Visser, S. P. Singlet versus triplet reactivity in an Mn(V)-Oxo species: Testing theoretical predictions against experimental evidence. *J. Am. Chem. Soc.* **2016**, *138*, 12375–12386.
- (58) Cantú Reinhard, F. G.; de Visser, S. P. Oxygen atom transfer using an iron(IV)-oxo embedded in a tetracyclic N-heterocyclic carbene system: How does the reactivity compare to Cytochrome P450 Compound I? *Chem. Eur. J.* **2017**, *23*, 2935–2944.
- (59) Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.
- (60) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B* **1988**, *37*, 785–789.
- (61) Hay, P. J.; Wadt, W. R. Ab initio effective core potentials for molecular calculations. Potentials for the transition metal atoms Sc to Hg. *J. Chem. Phys.* **1985**, *82*, 270–283.
- (62) Francl, M. M.; Pietro, W. J.; Hehre, W. J.; Binkley, J. S.; Gordon, M. S.; DeFrees, D. J.; Pople, J. A. Self-consistent molecular orbital methods. XXIII. A polarization-type basis set for second-row elements. *J. Chem. Phys.* **1982**, *77*, 3654–3658.
- (63) Tomasi, J.; Mennucci, B.; Cammi, R. Quantum mechanical continuum solvation models. *Chem. Rev.* **2005**, *105*, 2999–3093.
- (64) Cantú Reinhard, F. G.; Faponle, A. S.; de Visser, S. P. Substrate sulfoxidation by an iron(IV)-oxo complex: benchmarking computationally calculated barrier heights to experiment. *J. Phys. Chem. A* **2016**, *120*, 9805–9814.
- (65) Timmins, A.; Saint-André, M.; de Visser, S. P. Understanding how prolyl-4-hydroxylase structure steers a ferryl oxidant toward scission of a strong C–H bond. *J. Am. Chem. Soc.* **2017**, *139*, 9855–9866.
- (66) Cantú Reinhard, F. G.; Barman, P.; Mukherjee, G.; Kumar, J.; Kumar, D.; Kumar, D.; Sastri, C. V.; de Visser, S. P. Keto-enol tautomerization triggers an electrophilic aldehyde de-formylation reaction by a nonheme manganese(III)-peroxo complex. *J. Am. Chem. Soc.* **2017**, *139*, 18328–18338.
- (67) Quaroni, L. G.; Seward, H. E.; McLean, K. J.; Girvan, H. M.; Ost, T. W.; Noble, M. A.; Kelly, S. M.; Price, N. C.; Cheesman, M. R.; Smith, W. E.; Munro, A. W. Interaction of nitric oxide with cytochrome P450 BM3. *Biochemistry* **2004**, *43*, 16416–16431.
- (68) Munro, A. W.; Girvan, H. M.; Mason, A. E.; Dunford, A. J.; McLean, K. J. What makes a P450 tick? *Trends Biochem. Sci.* **2013**, *38*, 140–150.
- (69) Ouellet, H.; Lang, J.; Couture, M.; Ortiz de Montelano, P. R. Reaction of *Mycobacterium tuberculosis* cytochrome P450 enzymes with nitric oxide. *Biochemistry* **2009**, *48*, 863–872.
- (70) Luthra, A.; Denisov, I. G.; Sligar, S. G. Spectroscopic features of cytochrome P450 reaction intermediates. *Arch. Biochem. Biophys.* **2011**, *507*, 26–35.
- (71) Ost, T. W. B.; Clark, J.; Mowat, C. G.; Miles, C. S.; Walkinshaw, M. D.; Reid, G. A.; Chapman, S. K.; Daff, S. Oxygen activation and electron transfer in flavocytochrome P450 BM3. *J. Am. Chem. Soc.* **2003**, *125*, 15010–15020.
- (72) Sharma, S. K.; Schaefer, A. W.; Lim, H.; Matsumura, H.; Moënné-Loccoz, P.; Hedman, B.; Hodgson, K. O.; Solomon, E. I.; Karlin, K. D. Six-coordinate peroxynitrite low-spin iron(III) porphyrinate complex-the product of the reaction of nitrogen monoxide (·NO(g)) with a ferric-superoxide species. *J. Am. Chem. Soc.* **2017**, *139*, 17421–17430.
- (73) Liao, M.-S.; Huang, M.-J.; Watts, J. D. Binding of O₂ and NO to heme in heme-nitric oxide/oxygen-binding (H-NOX) proteins. A theoretical study. *J. Phys. Chem. B* **2013**, *117*, 10103–10114.
- (74) Green, M. T. Evidence for sulfur-based radicals in thiolate compound I intermediates. *J. Am. Chem. Soc.* **1999**, *121*, 7939–7940.
- (75) Shaik, S.; de Visser, S. P.; Ogliaro, F.; Schwarz, H.; Schröder, D. Two-state reactivity (TSR) mechanisms of hydroxylation and epoxidation by cytochrome P450 revealed by theory. *Curr. Opin. Chem. Biol.* **2002**, *6*, 556–567.
- (76) Shaik, S.; Kumar, D.; de Visser, S. P.; Altun, A.; Thiel, W. Theoretical perspective on the structure and mechanism of cytochrome P450 enzymes. *Chem. Rev.* **2005**, *105*, 2279–2328.
- (77) Ji, L.; Faponle, A. S.; Quesne, M. G.; Sainna, M. A.; Zhang, J.; Franke, A.; Kumar, D.; van Eldik, R.; Liu, W.; de Visser, S. P. Drug metabolism by cytochrome P450 enzymes: What distinguishes the pathways leading to substrate hydroxylation over desaturation? *Chem. Eur. J.* **2015**, *21*, 9083–9092.
- (78) Pickl, M.; Kurakin, S.; Cantú Reinhard, F. G.; Schmid, P.; Pöcheim, A.; Winkler, C. K.; Kroutil, W.; de Visser, S. P.; Faber, K.

Mechanistic studies of fatty acid activation by CYP₁₅₂ peroxygenases reveal unexpected desaturase activity. *ACS Catal.* **2019**, *9*, 565–577.

(79) de Visser, S. P.; Shaik, S. A proton-shuttle mechanism mediated by the porphyrin in benzene hydroxylation by cytochrome P450 enzymes. *J. Am. Chem. Soc.* **2003**, *125*, 7413–7424.

(80) Shaik, S.; Milko, P.; Schyman, P.; Usharani, D.; Chen, H. Trends in aromatic oxidation reactions catalyzed by cytochrome P450 enzymes: A valence bond modeling. *J. Chem. Theory Comput.* **2011**, *7*, 327–339.

(81) de Visser, S. P. Substitution of hydrogen by deuterium changes the regioselectivity of ethylbenzene hydroxylation by an oxo-iron-porphyrin catalyst. *Chem. Eur. J.* **2006**, *12*, 8168–8177.

(82) Balding, P. R.; Porro, C. S.; McLean, K. J.; Sutcliffe, M. J.; Maréchal, J.-D.; Munro, A. W.; de Visser, S. P. How do azoles inhibit cytochrome P450 enzymes? A density functional study. *J. Phys. Chem. A* **2008**, *112*, 12911–12918.

(83) Kishimoto, A.; Takagi, K.; Amano, A.; Sakurai, K.; Mizushima, T.; Shimada, H. Structure of a P450cam intermediate. Protein databank entry 3WRH. DOI: 10.2210/pdb3WRH/pdb.

(84) Berman, H. M.; Westbrook, J.; Feng, Z.; Gilliland, G.; Bhat, T. N.; Weissig, H.; Shindyalov, I. N.; Bourne, P. E. The Protein Data Bank. *Nucl. Acids Res.* **2000**, *28*, 235–242.

(85) Davydov, R.; Makris, T. M.; Kofman, V.; Werst, D. E.; Sligar, S. G.; Hoffman, B. M. Hydroxylation of camphor by reduced oxy-cytochrome P450cam: mechanistic implications of EPR and ENDOR studies of catalytic intermediates in native and mutant enzymes. *J. Am. Chem. Soc.* **2001**, *123*, 1403–1415.

(86) Kamachi, T.; Yoshizawa, K. A Theoretical study on the mechanism of camphor hydroxylation by compound I of cytochrome P450. *J. Am. Chem. Soc.* **2003**, *125*, 4652–4661.

(87) Schöneboom, S.; Cohen, S.; Lin, H.; Shaik, S.; Thiel, W. Quantum mechanical/molecular mechanical investigation of the mechanism of C-H hydroxylation of camphor by cytochrome P450cam: theory supports a two-state rebound mechanism. *J. Am. Chem. Soc.* **2004**, *126*, 4017–4034.

(88) Lai, R.; Li, H. Hydrogen abstraction of camphor catalyzed by cytochrome P450cam: A QM/MM Study. *J. Phys. Chem. B* **2016**, *120*, 12312–12320.

(89) Kumar, D.; Hirao, H.; de Visser, S. P.; Zheng, J.; Wang, D.; Thiel, W.; Shaik, S. New features in the catalytic cycle of cytochrome P450 during the formation of Compound I from Compound o. *J. Phys. Chem. B* **2005**, *109*, 19946–19951.

(90) de Visser, S. P. Trends in substrate hydroxylation reactions by heme and nonheme iron(IV)-oxo oxidants give correlations between intrinsic properties of the oxidant with barrier height. *J. Am. Chem. Soc.* **2010**, *132*, 1087–1097.

(91) Faponle, A. S.; Quesne, M. G.; de Visser, S. P. Origin of the regioselective fatty acid hydroxylation versus decarboxylation by a cytochrome P450 peroxygenase: What drives the reaction to biofuel production? *Chem. Eur. J.* **2016**, *22*, 5478–5483.

(92) Quesne, M. G.; Senthilnathan, D.; Singh, D.; Kumar, D.; Maldivi, P.; Sorokin, A. B.; de Visser, S. P. Origin of the enhanced reactivity of μ -nitrido-bridged diiron(IV)-oxo porphyrinoid complexes over cytochrome P450 Compound I. *ACS Catal.* **2016**, *6*, 2230–2243.

(93) Li, X.-X.; Postils, V.; Sun, W.; Faponle, A. S.; Solà, M.; Wang, Y.; Nam, W.; de Visser, S. P. Reactivity patterns of (protonated) Compound II and Compound I of Cytochrome P450: Which is the better oxidant? *Chem. Eur. J.* **2017**, *23*, 6406–6418.

(94) Liu, J. L.; Siegler, M. A.; Karlin, K. D.; Moënné-Loccoz, P. Direct resonance Raman characterization of a peroxynitrito copper complex generated from O₂ and NO and mechanistic insights into metal-mediated peroxynitrite decomposition. *Angew. Chem. Int. Ed.* **2019**, *58*, 10936–10940.

(95) Shaik, S.; de Visser, S. P.; Kumar, D. External electric field will control the selectivity of enzymatic-like bond activations. *J. Am. Chem. Soc.* **2004**, *126*, 11746–11749.

(96) Timmins, A.; Fowler, N. J.; Warwicker, J.; Straganz, G. D.; de Visser, S. P. Does substrate positioning affect the selectivity and reactivity in the hectochlorin biosynthesis halogenase? *Front. Chem.* **2018**, *6*, 513, doi: 10.3389/fchem.2018.00513.

(97) Wu, Y.; Fried, S. D.; Boxer, S. G. A Preorganized electric field leads to minimal geometrical reorientation in the catalytic reaction of ketosteroid isomerase. *J. Am. Chem. Soc.* **2020**, *142*, 9993–9998.

SYNOPSIS TOC:

